

JOURNAL OF ALGEBRA 134, 491–496 (1990)

A Remark on p -Radical Groups

SHIGEO KOSHITANI*

*Department of Mathematics, Faculty of Science,
Chiba University, Yayoi-cho, Chiba City, 260, Japan*

Communicated by Walter Feit

Received April 20, 1989

DEDICATED TO PROFESSOR KAZUHIKO HIRATA ON HIS 60TH BIRTHDAY

1. INTRODUCTION

Let FG be the group algebra of a finite group G over an algebraically closed field F of characteristic $p > 0$, and let P be a Sylow p -subgroup of G . We call G p -radical if the induced module $(F_P)^G$ from the trivial FP -module F_P is semi-simple (completely reducible) as a right FG -module. This definition is the same as that of Motose and Ninomiya in [7] (see [2, VI Corollary 6.4]).

The purpose of this note is to give one sufficient condition with respect to the Green correspondences under which G is p -radical. Namely,

THEOREM. *If the restriction $S_{N_G(Q)}$ of S to $F[N_G(Q)]$ is a simple $F[N_G(Q)]$ -module for any simple FG -module S where Q is a vertex of S , then G is p -radical.*

The condition in the theorem is not too strong, for the following reasons. Assume that G is p -radical. Let Q be any p -subgroup of G and let $N = N_G(Q)$. Then the theorems of Okuyama [8] and Alperin [1] say that the Green correspondence with respect to (G, Q, N) gives a bijection between the set of all non-isomorphic simple FG -modules with vertex Q and the set of all non-isomorphic simple FN -modules with vertex Q (see Proposition 1). Let S be a simple FG -module with vertex Q . Then the above fact means that the restriction S_N of S to FN is simple modulo a kind of projectives. Moreover, if Q is a Sylow p -subgroup of G , then S_N is

* Supported in part by the Alexander von Humboldt Foundation (West Germany) and Grants-in-Aid for Encouragement of Young Scientists (No. 61740012, No. 62740014, No. 63740012), the Ministry of Education, Science and Culture (Japan).

actually simple (see Proposition 2). Several years ago, Tsushima wrote a paper on p -radical groups [10]. The examples in his paper [10, Examples (1) and (2), p. 80] satisfy the condition in the theorem.

Throughout this paper, G is always a finite group and F is an algebraically closed field of characteristic $p > 0$, and we mean by an FG -module only a finitely generated right FG -module. We write $\dim V$ for the F -dimension of a vector space V over F . Let M and N be FG -modules. We write $\text{Ker } M$ for the kernel of M (see [2, p. 86]) and $\text{Soc } M$ for the socle of M as an FG -module. For a positive integer n , nM denotes the direct sum $M \oplus \cdots \oplus M$ (n times). We write $[M, N]_G$ for $\dim[\text{Hom}_{FG}(M, N)]$. We write $N|M$ if N is (isomorphic to) a direct summand of M . Let H be a subgroup of G and L an FH -module. Then L^G denotes the induced FG -module from L , say, $L^G = L \otimes_{FH} FH$, and M_H denotes the restriction of M to FH . We write F_G for the trivial FG -module. For ordinary characters χ and ψ of G , $(\chi, \psi)_G$ denotes the inner product in G . We write 1_G for the trivial ordinary character of G . We write $k(G)$ for the number of all conjugacy classes of G . The symbol \blacksquare means that a proof is finished. We write $O^{p'}(G)$ for the intersection of all normal subgroups of G of index prime to p . The notations $|G|$, $|G:H|$, H^x , $Z(G)$, $N_G(X)$, $C_G(X)$, $H \triangleleft G$, $\langle x \rangle$, and $O_{p',p}(G)$ are standard (see Gorenstein [3, p. 511]). Other notation and notions follow the books of Feit [2] and Landrock [6].

2. PROOF OF THE THEOREM

In order to prove our main result, we proceed by a sequence of several lemmas.

LEMMA 1. *Let $H \triangleleft G$ such that H is a p -group or p' -group. Assume that B and \bar{B} are p -blocks of G and G/H , respectively, such that $\bar{B} \subseteq B$ and that $Q \subseteq \text{Ker } S$ for any simple FG -module S in B where Q is a vertex of S . Then $V \subseteq \text{Ker } T$ for any simple $F(G/H)$ -module T in \bar{B} where V is a vertex of T .*

Proof. This follows from [4, Lemma 1.3(c)]. \blacksquare

LEMMA 2 (Knörr). *If a vertex of S is contained in $\text{Ker } S$ for any simple FG -module S in the principal p -block, then G is p -solvable.*

Proof. Let P be a Sylow p -subgroup of G . We may assume $P \neq 1$. Let S be any simple FG -module with vertex Q in the principal p -block. Then, by the theorem of Knörr [5, 3.7 Corollary (ii)], we may assume that $Z(P) \subseteq C_P(Q) \subseteq Q \subseteq P$. Hence, [2, IV Lemma 4.12(iii)] and the assumption imply that $1 \neq Z(P) \subseteq O_{p',p}(G)$. Thus, $G/O_{p',p}(G)$ is p -solvable from Lemma 1, [2, V Lemma 4.1], and induction. \blacksquare

LEMMA 3 (Okuyama). *Let S be a simple FG-module with vertex Q , and let fS be the Green correspondent of S with respect to $(G, Q, N_G(Q))$. Then fS is simple if and only if $S|(F_Q)^G$.*

Proof. Let $N = N_G(Q)$ and $T = fS$. Assume that T is simple. Then, by [2, III Corollary 4.13], T is a simple-projective $F(N/Q)$ -module. Hence $T|F(N/Q) \cong (F_Q)^N$, so that $S \cong f^{-1}T|T^G|(F_Q)^G$. The converse is proved by Okuyama [8, Lemma 2.2]. ■

LEMMA 4. *Let S be a simple FG-module with vertex Q , and let $N = N_G(Q)$. Then the following are equivalent.*

- (1) S_N is a simple FN-module.
- (2) S is a trivial source module and $Q \subseteq \text{Ker } S$.

Proof. (1) \Rightarrow (2): Easy by Lemma 3 and [2, III Corollary 2.13(i)].

(2) \Rightarrow (1): Let fS be the Green correspondent of S with respect to (G, Q, N) . Then $S_N = fS \oplus (\bigoplus Y_i)$ for indecomposable FN-modules Y_i such that Y_i is η -projective for all i where $\eta = \{A | A \text{ is a subgroup of } N \cap Q^x \text{ for some } x \in G - N\}$. There exist no such Y_i 's from [2, III Lemma 4.12] since $Q \subseteq \text{Ker } S$. Hence, we obtain (1) by Lemma 3. ■

LEMMA 5. *Let $H \triangleleft G$. If $S_{N_G(Q)}$ is simple for any simple FG-module S where Q is a vertex of S , then $T_{N_H(V)}$ is simple for any simple FH-module T where V is a vertex of T .*

Proof. Let T be a simple FH-module. Then, by Clifford theory and Frobenius reciprocity, $T|S_H$ for some simple FG-module S . Let Q be a vertex of S . By Lemma 4 and Mackey decomposition, T is a trivial source module and $V \subseteq \text{Ker } S$ for a vertex V of T . Hence, Lemma 4 implies the assertion. ■

The next lemma is a key of this paper. If G is p -radical, then G is p -solvable by Okuyama [9, Theorem 1]. Moreover, [7, Theorem 5] and inductive argument often make it possible to assume that G has a normal subgroup H of index p if we want to claim that G is p -radical. In general, G cannot be p -radical even if H is p -radical (see [7, Remark 2]). From this point of view, troubles come in when we claim that G would be p -radical.

LEMMA 6. *Let $H \triangleleft G$ and $|G:H| = p$. If H is p -radical and if $S_{N_G(Q)}$ is simple for any simple FG-module S where Q is a vertex of S , then G is also p -radical.*

Proof. Since G/H is cyclic and F is algebraically closed, we can assume from the theorem of Clifford that $\{S_i | i = 1, \dots, n\}$ and $\{T_j | j = 1, \dots, m\}$

$j = 1, \dots, p\} \cup \{T_i | i = m+1, m+2, \dots, n\}$ are the sets of all non-isomorphic simple FG - and FH -modules, respectively, for integers m and n with $m < n$ such that $(S_i)_H \cong \bigoplus_{j=1}^p T_{ij}$ and T_{i1}, \dots, T_{ip} are all conjugate in G for each $i \leq m$, and that $(S_i)_H \cong T_i$ for each $i \geq m+1$. Let J be the Jacobson radical of FG . By Frobenius reciprocity, $(T_{ij})^G \cong S_i$ for each $i \leq m$ and each j , and $\text{Soc}((T_i)^G) \cong (T_i)^G / (T_i)^G \cdot J \cong S_i$ for each $i \geq m+1$. Let P be a Sylow p -subgroup of G . Since H is p -radical, we can write

$$(F_{P \cap H})^H \cong \left(\bigoplus_{i \leq m} \bigoplus_j a_{ij} T_{ij} \right) \oplus \left(\bigoplus_{i \geq m+1} a_i T_i \right)$$

for positive integers a_{ij} and a_i .

Fix any $i \leq m$. By Frobenius reciprocity and Mackey decomposition, $[(F_P)^G, S_i]_G = [(F_P)^G, (T_{ij})^G]_G = [((F_P)^G)_H, T_{ij}]_H = [(F_{P \cap H})^H, T_{ij}]_H = a_{ij}$ for all j .

Next, fix any $i \geq m+1$. Let $S = S_i$, $T = S_H$, $a = a_i$, and $b = [(F_P)^G, S]_G$. By Lemma 4 and the assumption, S is a trivial source module. Clearly, T is also a trivial source module. Let (F, R, K) be a p -modular system (see [6, p. 47]). Recall that F is algebraically closed. Then S is liftable to an R -free trivial source RG -module \tilde{S} by [6, II Theorem 12.4(ii)]. Let χ be the K -character of G afforded by $\tilde{S} \otimes_R K$. Therefore, by Frobenius reciprocity and the result of Landrock [6, II Lemma 12.6(i)], $a = [(F_{P \cap H})^H, T]_H = [F_{P \cap H}, T_{P \cap H}]_{P \cap H} = \dim[\text{Soc}(T_{P \cap H})] = (\chi, 1_{P \cap H})_{P \cap H} = (1/|P \cap H|) \cdot \sum_{y \in P \cap H} \chi(y)$. Similarly, $b = (\chi, 1_P)_P = (1/|P|) \cdot \sum_{y \in P} \chi(y)$. From the assumption, Lemma 4, and [2, III Lemma 4.12], for each $y \in G$, y is in some vertex of S if and only if y is a p -element in $\text{Ker } S$. Let Y be the set of all such y 's. Take any $y \in Y$. Then $S_{\langle y \rangle} \cong (\dim S) \cdot F_{\langle y \rangle}$, so that $\chi(y) = \dim S$ by the result of Scott [6, II Lemma 12.6(ii)]. Moreover, $\chi(z) = 0$ for any $z \in P$ such that $z \notin Y$ by [6, II Lemma 12.6(iii)]. Therefore, $b = |P \cap \text{Ker } S| \cdot (\dim S) / |P|$. Similarly, we have $a = |P \cap H \cap \text{Ker } S| \cdot (\dim S) / |P \cap H|$. Let Q and V be vertices of S and T , respectively. Since G is p -solvable by the assumption and Lemmas 2 and 4, $|Q|/p = |V|$ by [4, Theorem 2.1]. Thus, $P \cap \text{Ker } S \neq P \cap H \cap \text{Ker } S$ by the assumption, Lemma 4, and [2, III Lemma 4.12]. Hence, $|P \cap \text{Ker } S| / |P \cap H \cap \text{Ker } S| = p$. Thus, $a = b$, so that $[(F_P)^G, S_i]_G = a_i$ for all $i \geq m+1$. This implies

$$(F_P)^G / (F_P)^G \cdot J \cong \left(\bigoplus_{i \leq m} a_{i1} S_i \right) \oplus \left(\bigoplus_{i \geq m+1} a_i S_i \right).$$

On the other hand, $\dim[(F_P)^G] = \dim[(F_{P \cap H})^H] = \sum_{i \leq m} a_{i1} (\sum_{j=1}^p \dim T_{ij}) + \sum_{i \geq m+1} a_i (\dim T_i) = \sum_{i \leq m} a_{i1} (\dim S_i) + \sum_{i \geq m+1} a_i (\dim S_i)$, which shows $(F_P)^G \cdot J = 0$. ■

Now, we have come to a position to prove our main result.

Proof of Theorem. If G has a normal subgroup of index p , then G is p -radical by Lemmas 5 and 6 and induction. So, we may assume $G \neq O^{p'}(G)$ since G is p -solvable by Lemmas 2 and 4. Hence, G is p -radical by Lemma 5, [7, Theorem 5], and induction. ■

EXAMPLES. The examples of p -radical groups mentioned in Tsushima's paper [10, Examples (1) and (2), p. 80], say, finite groups which have normal Sylow p -subgroups and the symmetric groups on 3 and 4 letters for all primes p , satisfy the condition in the Theorem.

3. PROPOSITIONS

In this section, we state two propositions and one corollary, which are mentioned in the introduction.

PROPOSITION 1 (Okuyama and Alperin). *Assume that G is p -radical. Let Q be any p -subgroup of G and let $N = N_G(Q)$. Then the Green correspondence with respect to (G, Q, N) gives a bijection between the set of all non-isomorphic simple FG-modules with vertex Q and the set of all non-isomorphic simple FN-modules with vertex Q .*

Proof. This follows from the properties of the Green correspondence [2, III Theorem 5.6], the theorems of Okuyama [8, Lemma 2.2] and Alperin [1, Lemma 1], [2, III Corollary 4.13], and the definition of p -radical groups. ■

PROPOSITION 2 (Okuyama). *Let P be a Sylow p -subgroup of G , and let $N = N_G(P)$. Assume that G is p -radical. Then we have the following.*

- (i) *If S is a simple FG-module such that P is a vertex of S , then S_N is a simple FN-module and $O^{p'}(G) \subseteq \text{Ker } S$.*
- (ii) *$k(G/O^{p'}(G)) = k(N/P)$.*

Proof. (i) follows from the result of Okuyama [9, Lemma 1], Lemma 4, and Sylow's theorem. (ii) is clear by (i), Proposition 1, and [2, III Corollary 2.13(i) and Lemma 4.12]. ■

COROLLARY. *Assume that G is p -radical and that $G = O^{p'}(G)$. Then we have the following.*

- (i) *If S is a simple FG-module whose vertex is a Sylow p -subgroup of G , then S is the trivial FG-module F_G .*
- (ii) *For a Sylow p -subgroup P of G , $N_G(P) = P$.*

Proof. (i) and (ii) follow from Proposition 2(i) and Proposition 2(ii), respectively. ■

Remark. In the corollary, the assumption that $G = O^{p'}(G)$ is not essential when we investigate the structure of G such that G is p -radical because of [7, Theorem 5] (see the remark just before Lemma 6).

ACKNOWLEDGMENTS

A part of this work was done while the author was at the Department of Mathematics, Essen University as a Humboldt research fellow (June–August 1988). The author expresses his great thanks to the Alexander von Humboldt Foundation for its financial support, and to Professor G. O. Michler and the people around him for their great hospitalities.

REFERENCES

1. J. L. ALPERIN, Weights for finite groups, in "The Arcata Conference on Representations of Finite Groups," Proceedings of Symposia in Pure Mathematics, Vol. 47, Part 1, 369–379, Amer. Math. Soc., Providence, 1987.
2. W. FEIT, "The Representation Theory of Finite Groups," North-Holland, Amsterdam, 1982.
3. D. GORENSTEIN, "Finite Groups," Chelsea, New York, 1980.
4. W. HAMERNIK AND G. MICHLER, On vertices of simple modules in p -solvable groups, *Mitt. Math. Sem. Giessen* **121** (1976), 147–162.
5. R. KNÖRR, On the vertices of irreducible modules, *Ann. of Math. (2)* **110** (1979), 487–499.
6. P. LANDROCK, "Finite Group Algebras and their Modules," London Mathematical Society Lecture Note Series, Vol. 84, Cambridge Univ. Press, Cambridge, U.K., 1983.
7. K. MOTOSE AND Y. NINOMIYA, On the subgroups H of a group G such that $J(KH)KG \supset J(KG)$, *Math. J. Okayama Univ.* **17** (1975), 171–176.
8. T. OKUYAMA, Module correspondence in finite groups, *Hokkaido Math. J.* **10** (1981), 299–318.
9. T. OKUYAMA, p -Radical groups are p -solvable, *Osaka J. Math.* **23** (1986), 467–469.
10. Y. TSUSHIMA, On p -radical groups, *J. Algebra* **103** (1986), 80–86.